

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

# **Long-Term Air Pollution and Traffic Noise Exposures and Mild Cognitive Impairment in Older Adults: A Cross-Sectional Analysis of the Heinz Nixdorf Recall Study**

Lilian Tzivian, Martha Dlugaj, Angela Winkler, Gudrun Weinmayr, Frauke Hennig, Kateryna B. Fuks, Mohammad Vossoughi, Tamara Schikowski, Christian Weimar, Raimund Erbel, Karl-Heinz Jöckel, Susanne Moebus, and Barbara Hoffmann, on behalf of the Heinz Nixdorf Recall Study Investigative Group

## **Table of Contents**

**Table S1.** Spearman correlations between air pollution (ESCAPE-LUR) and noise variables ( $r_s$ )

**Table S2.** Association of noise with MCI, OR (95% CI)

**Table S3.** Association between  $L_{DEN}$  (categorical) and overall MCI

**Figure S1.** Effect modification of association between exposures and overall MCI. Panel A: Associations of  $PM_{2.5}$  (per IQR) with overall MCI. Panel B: Associations of  $L_{DEN}$  (per 10 dB(A)) with overall MCI. Main model adjusted for age, sex, SES, alcohol consumption, smoking status, self-reported ETS, any regular physical activity and BMI.